

Absolute Total np and pp Cross Section Determinations

R.A. Arndt,¹ W.J. Briscoe,¹ A.B. Laptev,^{2,*} I.I. Strakovsky,¹ and R.L. Workman¹

¹*Center for Nuclear Studies, Department of Physics,
The George Washington University, Washington, DC 20052, USA*
²*Tulane University, New Orleans, LA 70118, USA*

PACS numbers: 11.80.Et,13.75.Cs,25.40.Cm,25.40.Dn

arXiv:0806.1198v2 [nucl-ex] 5 Jan 2009

*Present address: Los Alamos National Laboratory, Los Alamos, NM 87545, USA

Abstract — *Absolute total cross sections for np and pp scattering below 1000 MeV are determined based on partial-wave analyses of NN scattering data. These cross sections are compared with most recent ENDF/B and JENDL data files, and the Nijmegen partial-wave analysis. Systematic deviations from the ENDF/B and JENDL evaluations are found to exist in the low-energy region.*

I. INTRODUCTION

Nucleon-nucleon scattering is the simplest two-body reaction which allows an examination of different nuclear interaction models. Progress in the development of nuclear models is linked to the availability of high-quality data. np scattering is also used as a *primary* standard in measurements neutron-induced nuclear reactions [1, 2]. Its cross section is used in determining the flux of incoming neutrons.

This information is important in many applications, such as astrophysics [(n, γ), (n, α) and others] and the transmutation of nuclear waste [(n, f), (n, γ) and others], energy generation and the conceptual design of an innovative nuclear reactor being carried out in the course of the Generation IV initiative [3] [(n, f), and neutron-actinoid elastic and inelastic scattering and others]. The increasing quality of neutron-induced nuclear reaction measurements requires a high-quality standard for np cross sections, reproducing total np cross sections with an accuracy of 1% or better for energies below 20 MeV [1, 4]. The need for neutron data above 20 MeV up to hundreds of MeV with accuracy better than 10% [4] leads to the requirement of cross section data for the np *reference* reaction with uncertainties at the few percent level.

An extensive database exists for nucleon-nucleon scattering, with measurements from laboratories worldwide. These datasets, from the various laboratories, have different statistical and systematical uncertainties which must be taken into account when combined into a single fit. At present, there are several evaluations of total np cross sections below 20 MeV. Perhaps most widely known are the ENDF/B [5] and JENDL [6] nuclear data files. An R-matrix analysis of the NN system [7] was used in course of the ENDF/B evaluation of np cross sections whereas, in the JENDL np total cross section evaluation, a method based on phase-shift data [8, 9] was used.

Here we will concentrate on total np and pp cross section determined on the basis of

recent energy-dependent (global) fits and associated single-energy solutions (SES) from the George Washington (GW) Data Analysis Center [10]. Precise measurements collected over many years have helped to isolate discrepancies between different experiments and have contributed to a good description of nucleon-nucleon scattering at the level of both observables and amplitudes.

In Section II, we comment on the np and pp data which are available in the GW database and which have been used in this analysis. In Section III, we give a brief overview of the method used to fit data and extract amplitudes. Then, in Section IV, we present total np and pp cross sections determined on the basis of both global and SES results. Finally, in Section V, we summarize our findings.

II. DATABASE

The GW fit to NN elastic scattering data covers an energy range from threshold up to a laboratory kinetic energy of 1300 MeV (for np data) and 3000 MeV (for pp data). The np analysis was restricted to 1300 MeV due to a lack of high-energy data. The full database includes all available unpolarized and polarized measurements. A number of fits, from the GW group and others, are available through the on-line SAID facility [11].

The evolution of the SAID database is summarized in Table I. Over the course of seven previous GW NN partial-wave analyses (PWA) [8, 9, 10, 12, 13, 14, 15], the upper energy limit has been extended from 1200 (1100) to 3000 (1300) MeV in the laboratory proton (neutron) kinetic energy. (These fits are regularly updated online, and the designation SM97, for example, denotes the time of update - summer 1997.)

Not all of the available data have been used in each fit. Some data with very large χ^2 contributions have been excluded. Redundant data are also excluded. These include total elastic cross sections based on differential cross sections already contained in the database. Polarized measurements with uncertainties greater than 0.2 are not included as they have little influence in GW fits. However, all available data have been retained in the database (the excluded data labeled as “flagged”) so that comparisons can be made through the GW on-line facility [11]. A complete description of the database, and those data not included in GW fits, is available from the authors [11].

III. PARTIAL-WAVE ANALYSIS

Simultaneous fits to the full database are possible within the formalism used and described in previous GW analyses [8, 13, 14]. The observables are represented in terms of partial-wave amplitudes, using a Chew-Mandelstam K-matrix approach, which incorporates the effect of an $N\Delta$ channel on the NN scattering process. By parameterizing the K-matrix elements as functions of energy, data from threshold to 3000 MeV can be fitted simultaneously (both pp and np , with a 1300 MeV limit for np). In general, GW partial-wave analyses have attempted to remain as model-independent as possible. The present (SP07) and previous energy-dependent solutions are compared in Table I.

In fitting the data, systematic uncertainty has been used as an overall normalization factor for angular distributions. With each angular distribution, we associate the pair (X, ϵ_X) : a normalization constant (X) and its uncertainty (ϵ_X). The quantity ϵ_X is generally associated with the systematic uncertainty (if known). The modified χ^2 function, to be minimized, is then given by

$$\chi^2 = \sum_i \left(\frac{X\theta_i - \theta_i^{\text{exp}}}{\epsilon_i} \right)^2 + \left(\frac{X - 1}{\epsilon_X} \right)^2,$$

where the subscript i labels data points within the distribution, θ_i^{exp} is an individual measurement, θ_i is the calculated value, and ϵ_i is the statistical uncertainty. For total cross sections and excitation data, we have combined statistical and systematic uncertainties in quadrature. Renormalization freedom significantly improves GW best-fit results, as shown in Table II.

Starting from this global fit, we have also generated a series of SES results. Each SES is based on a 'bin' of scattering data spanning a narrow energy range. A total 43 SES have been generated, with central energy values ranging from 5 MeV to 2830 MeV, and bin widths varying from 2 to 75 MeV. For example, solution c5 is a fit to data between 4 and 6 MeV. In generating the SES, a linearized energy dependence is taken over the energy range, reducing the number of searched parameters. A systematic deviation between the SES and global fits is an indication of missing structure in the global fit (or possibly problems with a particular dataset). An error matrix is generated in the SES fits, which can be used to estimate the overall uncertainty in the global fit. The global and SES fit results, up to 1000 MeV, are summarized in Table III (further details are given in Ref. [10]).

IV. TOTAL np AND pp CROSS SECTIONS

Isovector and isoscalar partial-wave amplitudes, determined through the partial-wave analysis, have been used to generate total np and pp cross sections presented in Tables IV and V. In Table IV, we compare results from two global fits, SP07 (with an energy limit of 3000 MeV) and LE08 (a low-energy fit to 25 MeV which searches 19 parameters, scattering length a and effective range r for 3 S-waves and 13 leading parameters for S, P, and D waves). LE08 results in a $\chi^2/\text{data} = 696/391$ for pp and $627/631$ for np . The SP07, LE08, and SES results below 20 MeV, are summarized in Table VI. Errors for LE08 have been generated from the error matrix and require some comments.

In the region below 25 MeV, there are numerous total cross section measurements for np , but not for pp which is hindered by large Coulomb effects. As a result, the np error estimates are more reasonable. Those quoted for pp are far too small (lower limits) in the threshold region. In two cases (c5 and c150 of Table III), the χ^2 from the energy-dependent fit was actually lower than that from the SES fit. There is a possibility that this can indicate that the energy bin for the SES fit is too broad. But a rather narrow bin will not allow a stable result because of the lack of data to constrain the solution.

For the region above 25 MeV, in Table V, we compare the global fit SP07 with the grid of SES fits. Here the SES errors give a more accurate estimate of the uncertainty in our cross sections. The amplitudes found in GW fits to 1000 MeV have remained stable against the addition of new measurements for many years. Sufficient observables exist for a direct amplitude reconstruction at many energies, and we have compared GW amplitudes to those found in this way in Ref. [10].

As cross sections change rapidly near threshold, we have chosen to display the agreement between various fits in terms of ratios. This gives a clear picture of the overall consistency and reveals cases where systematic deviations are present. The ratios of SES values to the global fit SP07, below 20 MeV, are displayed in Fig. 1(a). Also plotted is a band showing the ratio of LE08 to SP07 determinations of the np cross section. As expected, this band more closely reproduces the np SES, plotted as single points with error bars, than the 3000 MeV fit SP07. Deviations are within 1% for the np determinations, and within 2% for pp .

In Fig. 1(b), we plot ratios of SP07 and SES, for both np and pp cases, to the Nijmegen PWA predictions [16]. The low-energy Nijmegen total pp cross sections are systematically

above SP07 (about 2% or less) while np cross sections agree with SP07 at better than the 0.3% level.

In Fig. 2, we plot ratios of the GW np fits with the ENDF/B and JENDL nuclear data files [5, 6]. A slightly better agreement is found with JENDL [6] than with ENDF/B [5], though the wiggles seen in Fig. 2(b) reflect a lack of smoothness in JENDL (SP07 and LE08 are a smooth function of energy). The ENDF/B result is systematically below SP07 and the Nijmegen fit [16], but the maximal deviation is only 1%. SP07 and JENDL agree at the 0.5% level over most of the region below 20 MeV.

At higher energies (up to 1000 MeV), ratios of the grid of SES to SP07 differ from unity by less than 3% [Fig. 3(a)]. Above 180 MeV, SAID np cross sections are larger than JENDL/HE [6] by up to 5% [Fig. 3(b)].

V. SUMMARY AND CONCLUSIONS

We have generated fits to describe the total np and pp scattering cross sections below 1000 MeV. These fits have been both energy dependent (SP07, LE08) and single-energy (analyzing narrow bins of data). The uncertainties associated with our total np cross sections below 20 MeV are clearly less than 1%. The agreement between SP07, JENDL, and the Nijmegen analysis, suggests an uncertainty of 0.5% or less. A comparison with ENDF/B shows deviations of 1% or less. Errors on the LE08 solution in Table IV, while obtained using a well-defined method, are lower bounds as they do not account for systematics effects.

For the pp cross sections, uncertainties are larger and systematic disagreements are evident in comparisons with the Nijmegen PWA. The main problem stems from a lack of relevant pp data at low energies. Here also, at low energies, the various determinations agree at the few-percent level.

The advantage of the GW parameterization is its smooth energy dependence and coverage from threshold to high energies. We also have the capability to modify the GW fits to either generate SES centered on a particular energy, or produce lower-energy fits when a specific energy region is of interest. We will continue to update both GW energy-dependent and single-energy solutions as the new measurements become available.

Acknowledgments

The authors express their gratitude to A.D. Carlson, G.M. Hale, and T. Hill for the useful discussion. This work was supported in part by the U.S. Department of Energy under Grant DE-FG02-99ER41110. The authors acknowledge partial support from Jefferson Lab, by the Southeastern Universities Research Association under DOE contract DE-AC05-84ER40150.

-
- [1] S. BADIKOV, CHEN ZHENPENG, A.D. CARLSON *et al.*, “International Evaluation of Neutron Cross-Section Standards,” Technical Report STI/PUB/1291 (International Atomic Agency, Vienna, 2007).
 - [2] *Proc. of the CSEWG and USNDP Meetings, BNL, Nov. 2007*, P. OBLOZINSKY, Ed.; <http://www.nndc.bnl.gov/meetings/csewg2007/>
 - [3] “A Technology Roadmap for Generation IV Nuclear Energy Systems,” by U.S. DOE Nuclear Energy Research Advisory Committee and the Generation IV International Forum, Dec. 2002, Report GIF-002-00; <http://www.ne.doe.gov/geniv/documents/genivroadmap.pdf>.
 - [4] G. ALBBERTI, G. PALMIOTTI, M. SALVATOIRES *et al.*, “Nuclear data sensitivity, uncertainty and target accuracy assessment for future nuclear systems,” *Ann. Nucl. Energy*, **33**, 700 (2006).
 - [5] M. B. CHADWICK, P. OBLOZINSKY, M. HERMAN *et al.*, “ENDF/B-VII.0: Next generation evaluated nuclear data library for nuclear science and technology,” *Nucl. Data Sheets*, **107**, 2931 (2006); numerical data are available via National Nuclear Data Center (NNDC); <http://www.nndc.bnl.gov/endl>, release ENDF/B-VII.0.
 - [6] K. SHIBATA, T. KAWANO, T. NAKAGAWA *et al.*, “Japanese evaluated nuclear data library version 3 revision-3: JENDL-3.3,” *J. Nucl. Sci. Tech.* **39**, 1125 (2002); numerical data are available via JAEA Nuclear Data Center; <http://www.ndc.jaea.go.jp/jendl/j33/j33.html>, release JENDL-3.3.
 - [7] G. M. HALE and A. S. JOHNSON, “Results for n+p capture from an R-matrix analysis of N-N scattering,” in *Proc. 17th Int. IUPAP Conf. Few-Body Problems in Physics (Few-Body 17)*, June 2003, Durham, NC, U.S.A., W. GLOECKLE and W. TORNOW, Eds. (Elsevier B.V., 2004) p. S120.
 - [8] R. A. ARNDT, J. S. HYSLOP, and L. D. ROPER, “Nucleon-nucleon partial-wave analysis to 1100 MeV,” *Phys. Rev. D* **35**, 128 (1987).
 - [9] R. A. ARNDT, L. D. ROPER, R. A. BRYAN, R. B. CLARK, B. J. VER-WEST, and P. SIGNELL, “Nucleon-nucleon partial-wave analysis to 1 GeV,” *Phys. Rev. D* **28**, 97 (1983).
 - [10] R. A. ARNDT, W. J. BRISCOE, I. I. STRAKOVSKY, and R. L. WORKMAN, “Updated analysis of NN elastic scattering to 3 GeV,” *Phys. Rev. C* **76**, 025209 (2007); arXiv:0706.2195

[nucl-th].

- [11] The full database and numerous PWAs with potential models can be accessed via ssh call to the SAID facility `gwdac.phys.gwu.edu`, with userid: `said` (no password), or a link to the website: <http://gwdac.phys.gwu.edu>.
- [12] R. A. ARNDT, I. I. STRAKOVSKY, and R. L. WORKMAN, “Nucleon-nucleon elastic scattering to 3 GeV,” *Phys. Rev. C* **62**, 034005 (2000); [nucl-th/0004039].
- [13] R. A. ARNDT, C. H. OH, I. I. STRAKOVSKY, R. L. WORKMAN, and F. DOHRMANN, “Nucleon-nucleon elastic scattering analysis to 2.5 GeV,” *Phys. Rev. C* **56**, 3005 (1997); [nucl-th/9706003].
- [14] R. A. ARNDT, I. I. STRAKOVSKY, and R. L. WORKMAN, “Updated analysis of NN elastic scattering data to 1.6 GeV,” *Phys. Rev. C* **50**, 2731 (1994); [nucl-th/9407035].
- [15] R. A. ARNDT, L. D. ROPER, R. L. WORKMAN, and M. W. McNAUGHTON, “Nucleon-nucleon partial-wave analysis to 1.6 GeV,” *Phys. Rev. D* **45**, 3995 (1992).
- [16] V. G. J. STOKS, R. A. M. KLOMP, M. C. M. RENTMEESTER, and J. J. de SWART, “Partial-wave analysis of all nucleon-nucleon scattering data below 350 MeV,” *Phys. Rev. C* **48**, 792 (1993); <http://nn-online.org/>.

TABLE I: Comparison of recent SP07 [10] and previous SP00 [12], SM97 [13], SM94 [14], FA91 [15], SM86 [8], and SP82 [9] GW energy-dependent partial-wave analyses. The χ^2 values for previous solutions correspond to published results.

Solution	Range (MeV)	χ^2/np data	Range (MeV)	χ^2/pp data
SP07	0–1300	21496/12693	0–3000	44463/24916
SP00	0–1300	18693/11472	0–3000	36617/21796
SM97	0–1300	17437/10854	0–2500	28686/16994
SM94	0–1300	17516/10918	0–1600	22371/12838
FA91	0–1100	13711/ 7572	0–1600	20600/11880
SM86	0–1100	8871/ 5474	0–1200	11900/ 7223
SP82	0–1100	9103/ 5283	0–1200	9199/ 5207

TABLE II: χ^2/data (pp and np) normalized (Norm) and unnormalized (Unnorm) below 1000 MeV for GW fits SP07 [10], SP00 [12], SM97 [13], and the Nijmegen PWA fit [16] (the Nijmegen solution is valid up to 350 MeV - the final tabulated energy range is 200–350 MeV). Last two columns list numbers of np and pp data included in fit SP07 and its associated SES.

Range (MeV)	SP07 Norm/Unnorm	SP00 Norm/Unnorm	SM97 Norm/Unnorm	Nijmegen93 Norm/Unnorm	Data np	Data pp
0– 4	2.5 / 28.0	2.5 / 28.0	2.5 / 27.9	3.3 / 26.7	63	193
0– 20	1.8 / 13.2	1.9 / 13.3	2.3 / 13.8	2.9 / 10.1	468	389
0– 200	1.5 / 7.2	1.5 / 7.1	1.7 / 6.8	1.7 / 5.9	2381	1491
200– 400	1.3 / 3.3	1.3 / 3.3	1.4 / 3.3	1.3 / 2.5	2208	2172
400– 600	1.5 / 8.7	1.4 / 8.0	1.5 / 10.6		2779	3635
600– 800	1.5 / 8.4	1.5 / 7.7	1.4 / 10.6		2529	3974
800–1000	1.4 / 3.4	1.4 / 3.4	1.4 / 3.1		2112	3274

TABLE III: Single-energy (binned) fits [10] of combined elastic np and pp scattering data below 1000 MeV. χ_E^2 is given by the energy-dependent fit, SP07 [10], over the same energy intervals.

Solution	Range (MeV)	pp χ_E^2	χ^2/data	np χ_E^2	χ^2/data
c5	4– 6	37	22/ 28	74	77/ 62
c10	7– 12	128	81/ 88	255	222/106
c15	11– 19	55	16/ 27	366	219/247
c25	19– 30	251	120/ 114	319	293/316
c50	32– 67	331	283/ 224	721	660/467
c75	60– 90	61	48/ 72	652	516/355
c100	80– 120	155	149/ 154	521	437/389
c150	125– 174	352	311/ 287	533	542/352
c200	175– 225	570	528/ 435	870	742/519
c250	225– 270	289	266/ 263	601	543/438
c300	276– 325	853	815/ 805	1163	1074/770
c350	325– 375	522	503/ 474	509	462/381
c400	375– 425	819	771/ 680	1295	1208/805
c450	425– 475	1307	1186/ 877	1216	1194/912
c500	475– 525	1811	1556/1215	1386	1240/815
c550	525– 575	1075	938/ 817	1192	987/719
c600	575– 625	1196	1045/ 838	517	423/367
c650	625– 675	980	959/ 807	1502	1265/877
c700	675– 725	982	956/ 887	460	396/386
c750	725– 775	1474	1085/ 926	545	508/382
c800	775– 824	2056	1833/1385	1777	1427/950
c850	827– 874	1296	1142/ 980	500	393/365
c900	876– 924	542	406/ 444	936	730/627
c950	926– 974	961	769/ 728	528	356/353
c999	976–1020	1206	1010/ 776	421	274/329

TABLE IV: Comparison of SP07 [10] and LE08 results for total np (σ^n) and pp (σ^p) cross sections below 25 MeV. See text for a discussion of uncertainties.

T (MeV)	LE08 σ^p (mb)	SP07 σ^p (mb)	LE08 σ^n (mb)	SP07 σ^n (mb)	T (MeV)	LE08 σ^p (mb)	SP07 σ^p (mb)	LE08 σ^n (mb)	SP07 σ^n (mb)
0.5	1100.00±0.01	1098.00	6148.00±0.07	6135.00	13.0	264.00±0.10	263.70	745.40±0.12	745.00
1.0	1513.00±0.01	1513.00	4261.00±0.19	4253.00	13.5	253.60±0.11	253.00	719.10±0.13	718.80
1.5	1469.00±0.01	1471.00	3421.00±0.32	3417.00	14.0	243.90±0.13	243.10	694.40±0.14	694.20
2.0	1323.00±0.01	1325.00	2911.00±0.44	2911.00	14.5	234.90±0.14	233.80	671.10±0.14	671.00
2.5	1172.00±0.01	1175.00	2555.00±0.52	2558.00	15.0	226.60±0.15	225.20	649.00±0.15	649.20
3.0	1040.00±0.01	1042.00	2286.00±0.57	2291.00	15.5	218.80±0.16	217.10	628.20±0.16	628.60
3.5	927.70±0.01	930.60	2073.00±0.61	2079.00	16.0	211.60±0.17	209.60	608.40±0.17	609.10
4.0	833.90±0.01	836.70	1898.00±0.61	1906.00	16.5	204.80±0.18	202.50	589.70±0.18	590.60
4.5	755.00±0.01	757.70	1752.00±0.61	1760.00	17.0	198.50±0.20	195.80	571.90±0.19	573.10
5.0	688.10±0.01	690.60	1627.00±0.59	1635.00	17.5	192.60±0.21	189.50	555.00±0.20	556.40
5.5	630.90±0.01	633.30	1519.00±0.55	1526.00	18.0	187.00±0.23	183.60	539.00±0.21	540.60
6.0	581.60±0.02	583.80	1424.00±0.52	1431.00	18.5	181.80±0.24	178.00	523.70±0.23	525.50
6.5	538.70±0.02	540.80	1341.00±0.47	1347.00	19.0	176.90±0.25	172.80	509.20±0.24	511.20
7.0	501.10±0.02	503.10	1266.00±0.43	1271.00	19.5	172.30±0.27	167.80	495.40±0.25	497.50
7.5	468.00±0.03	469.90	1199.00±0.37	1203.00	20.0	167.90±0.28	163.00	482.30±0.26	484.40
8.0	438.70±0.03	440.40	1139.00±0.33	1142.00	20.5	164.00±0.30	158.70	469.80±0.28	471.90
8.5	412.50±0.04	414.00	1084.00±0.29	1086.00	21.0	160.20±0.32	154.40	457.90±0.29	460.00
9.0	389.00±0.04	390.30	1033.00±0.25	1035.00	21.5	156.50±0.33	150.40	446.50±0.31	448.50
9.5	367.80±0.05	369.00	987.10±0.21	988.60	22.0	153.10±0.35	146.50	435.70±0.32	437.60
10.0	348.60±0.06	349.60	944.60±0.18	945.50	22.5	149.90±0.36	142.80	425.30±0.34	427.00
10.5	331.20±0.06	332.00	905.30±0.15	905.70	23.0	146.80±0.38	139.30	415.50±0.36	417.00
11.0	315.40±0.07	316.00	868.80±0.13	868.90	23.5	143.90±0.40	135.90	406.10±0.37	407.30
11.5	300.90±0.08	301.20	834.80±0.12	834.60	24.0	141.20±0.41	132.70	397.10±0.39	398.00
12.0	287.50±0.09	287.70	803.00±0.12	802.70	24.5	138.60±0.43	129.60	388.50±0.41	389.00
12.5	275.30±0.10	275.20	773.30±0.12	772.90	25.0	136.20±0.45	126.70	380.30±0.43	380.40

TABLE V: Comparison of SES and SP07 [10] results for total np (σ^n) and pp (σ^p) cross sections between 25 and 1000 MeV.

T (MeV)	Solution	SES σ^p (mb)	SP07 σ^p (mb)	SES σ^n (mb)	SP07 σ^n (mb)
25.0	c25	128.20 $\pm$ 0.43	126.70	380.90 $\pm$ 0.63	380.40
50.0	c50	58.78 $\pm$ 0.12	58.95	168.10 $\pm$ 0.35	168.40
75.0	c75	39.14 $\pm$ 0.27	39.80	101.70 $\pm$ 0.32	103.70
100.0	c100	31.11 $\pm$ 0.30	31.70	73.62 $\pm$ 0.35	75.53
150.0	c150	25.40 $\pm$ 0.09	25.74	50.77 $\pm$ 0.31	52.24
200.0	c200	23.66 $\pm$ 0.12	24.22	42.43 $\pm$ 0.22	43.04
250.0	c250	23.89 $\pm$ 0.19	23.89	37.89 $\pm$ 0.36	38.35
300.0	c300	23.46 $\pm$ 0.13	24.07	35.53 $\pm$ 0.21	35.61
350.0	c350	24.53 $\pm$ 0.13	24.93	34.05 $\pm$ 0.31	34.11
400.0	c400	25.91 $\pm$ 0.09	26.17	33.70 $\pm$ 0.17	33.34
450.0	c450	28.53 $\pm$ 0.14	28.06	34.13 $\pm$ 0.18	33.24
500.0	c500	30.60 $\pm$ 0.18	30.83	34.30 $\pm$ 0.23	33.82
550.0	c550	34.13 $\pm$ 0.32	34.42	35.24 $\pm$ 0.26	34.97
600.0	c600	37.29 $\pm$ 0.33	38.13	35.32 $\pm$ 0.13	36.30
650.0	c650	40.54 $\pm$ 0.36	41.30	37.86 $\pm$ 0.29	37.43
700.0	c700	43.53 $\pm$ 0.49	43.65	37.40 $\pm$ 0.96	38.23
750.0	c750	46.23 $\pm$ 0.31	45.12	38.38 $\pm$ 0.94	38.66
800.0	c800	45.83 $\pm$ 0.24	45.88	39.15 $\pm$ 0.21	38.80
850.0	c850	45.97 $\pm$ 0.56	46.20	37.71 $\pm$ 0.49	38.80
900.0	c900	47.32 $\pm$ 0.82	46.30	38.27 $\pm$ 0.59	38.76
950.0	c950	47.49 $\pm$ 0.62	46.30	37.69 $\pm$ 0.64	38.74
1000.0	c999	47.19 $\pm$ 0.64	46.25	38.14 $\pm$ 0.57	38.78

TABLE VI: Single-energy (binned) fits [10] of combined elastic np and pp scattering data below 20 MeV. χ_E^2 are given by the energy-dependent SP07 [10], LE08, and NI93 [16] fits, over the same energy intervals. Number of searched parameters in SES is given in the 2nd column in brackets.

Solution	Range (MeV)	SP07 χ_E^2	LE08 χ_E^2	SES χ^2	NI93 χ_E^2/data
for pp					
c5	4 – 6 (6)	37	30	22	51/28
c10	7 – 12 (6)	129	98	81	107/88
c15	11 – 19 (8)	55	37	17	15/27
for np					
c5	4 – 6 (6)	75	58	78	78/ 62
c10	7 – 12 (6)	256	138	222	132/106
c15	11 – 19 (8)	366	231	219	246/247




FIG. 1: Ratios of total np and pp cross sections below 20 MeV. Horizontal bars give the energy binning of SES (Table III). (a) Single-energy to energy-dependent SP07 [10] ratios are plotted. np (pp) results are shown as solid (open) circles. The band represents the ratio of LE08 to SP07 for the np case. (b) SES (solid circles for np and open circles for pp) and SP07 (solid line for np and dash-dotted line for pp) divided by Nijmegen PWA predictions [16] are plotted. The band represents the ratio of LE08 to Nijmegen PWA for the np case.




FIG. 2: Ratio of total np cross sections below 20 MeV. Horizontal bars give the energy binning of SES (Table III). (a) Single-energy (solid circles) and SP07 (solid line) fits [10] divided by the ENDF/B [5] results are plotted. The band gives a ratio of LE08 to ENDF/B. (b) The same for JENDL [6] evaluated data. The wiggles seen in Fig. 2(b) reflect a lack of smoothness in JENDL (SP07 and LE08 are a smooth function of energy).




FIG. 3: Ratios of total np and pp cross sections between 20 and 1000 MeV. (a) SES to SP07 [10] ratios are plotted. np (pp) results are shown as solid (open) circles. (b) np SES (solid circles) and SP07 (solid line) divided by JENDL/HE [6] results are plotted.